

Transition linewidth of cross-correlations in random intensity fluctuations in EIT

Lei Feng,¹ Pengxiong Li,¹ Tun Wang,^{2,3} and Yanhong Xiao¹

¹*Department of Physics, State Key Laboratory of Surface Physics, Laboratory of Advanced Materials and Key Laboratory of Micro and Nano Photonic Structures (Ministry of Education), Fudan University, Shanghai 200433, China*

²*Department of Physics, University of Connecticut, Storrs, Connecticut 06269, USA*

³*Institute of Care-life, Chengdu 610041, China*

(Dated: March 8, 2013)

It is known that cross-correlation between the random intensity fluctuations of two lasers forming electromagnetically-induced transparency (EIT) exhibits a transition from correlation to anticorrelation. We study the linewidth behavior of this transition and have found the linewidth is below the (effective) coherence lifetime limit and is only limited by competing noises. We established a numerical model which reveals the linewidth dependence on laser linewidth and laser power. Our experiments using lasers with different linewidth showed results in qualitative agreement with the model. This result is useful for quantum optics using EIT, and may also have applications in spectroscopy and precision measurements.

PACS numbers: 32.70.Jz, 42.50.Gy, 42.62.Fi

Manipulating photon statistics and laser noise through coherent atom-light interaction has recently become a topic of great interest for quantum optics, spectroscopy and quantum information processing. Electromagnetically induced transparency (EIT) [1] or coherent population trapping (CPT) [2] is one of the most widely used processes for such purposes. Interesting findings on both the quantum fluctuations and the classical noises in EIT/CPT fields have been reported. For instance, a CPT system can be used to generate entangled bright photon pairs [3] and spin squeezing simultaneously [4]; EIT is opaque to quantum fluctuations in laser fields [5]; EIT-like features exist in the noise spectrum of lasers [6–8] even when far off EIT resonance; a cross-correlation resonance [9] in an EIT system can have linewidth below the coherence lifetime limit [10] by using frequency-modulated laser fields.

In this paper, we study the cross-correlation resonance in the random noise case, especially for lasers with diffusive phase noises. It has been known that the cross-correlation between the random intensity fluctuations (converted from the random phase noise in lasers) in the two EIT optical fields exhibits a transition from correlation to anticorrelation when the Raman (two-photon) detuning is varied, with a linewidth narrower than that of EIT [8, 9]. However, it remains unclear what determines the linewidth. Here, we report the finding that the linewidth of the transition depends on the laser linewidth and laser power, and is below the coherence lifetime limit. In principle the linewidth can approach zero for infinitely narrow laser width if other noise sources are neglected. We established a numerical model which predicts linewidth behaviors in good qualitative agreement with our experimental results. Although our current study is on classical noise, the result is meaningful for quantum noise experiments since laser phase noise is inevitable and laser phase noise to intensity noise conversion (PN-IN) [11] may overwhelm quantum correlations.

Meanwhile, this study can contribute to the interesting field of noise spectroscopy [12, 13], and can also be useful for precision measurement purpose since such PN-IN process degrades performance of atomic frequency standards [6].

The physics picture for the sharp transition from correlation to anticorrelation was provided in [10] for frequency-modulated lasers. It is still applicable here since the random laser noise can be viewed as the sum of frequency modulation at all frequencies [14]. We consider a typical CPT configuration (see Fig. 1) where the pump E_1 and probe E_2 have equal Rabi frequencies and identical phase noises originated from the same laser source. They couple the excited state $|1\rangle$ to two ground states $|2\rangle$, $|3\rangle$ respectively. Although the physics is valid for a generic Λ system, we consider the Zeeman EIT for simplicity. When the two-photon detuning $\Delta = 0$, E_1 and E_2 have the same one-photon detuning, and hence experience parallel PN-IN slopes leading to correlations; when $\Delta \neq 0$, the sharp dispersion associated with the ground state coherence breaks the symmetry and induces a large offset ($\gg |\Delta|$) between the minima of the two transmission spectra [10]. This offset gives rise to opposite PN-IN slopes around $\delta = 0$ resulting in anticorrelated IN [15]. This picture indicates that narrower laser linewidth leads to sharper transition from correlation to anticorrelation, as shall be verified by our theory and experiments below.

In our model, we assume the Rabi frequencies for E_1 and E_2 are $\Omega_1 = \Omega_2 = \Omega e^{i\phi(t)}$, with $\phi(t)$ being the random phase noise obeying Gaussian statistics, known as Wiener-Levy process [16, 17]:

$$\langle \dot{\phi}(t) \rangle = 0, \langle \dot{\phi}(t) \dot{\phi}(t') \rangle = 2D\delta(t - t') \quad (1)$$

where D is the half width of laser spectrum at half maximum (HWHM). As in Eq. (1), the phase changing rate $\dot{\phi}(t)$ does not depend on its historical fluctuations. Dynamics of the density matrix is determined by




FIG. 1: (Color online) Intuitive picture for the sharp transition from correlation to anticorrelation in the intensity fluctuations of the two EIT fields E_1 and E_2 . δ is the averaged one-photon detuning, and Δ is the two-photon detuning, generated by shifting the energy levels with a magnetic field via Zeeman shift. (See text for details).

the stochastic optical Bloch equation [18] and the Hamiltonian is:

$$\mathbf{H} = \Omega_1|3\rangle\langle 1| + \Omega_2|3\rangle\langle 2| - \delta|3\rangle\langle 3| + \Delta/2(|2\rangle\langle 2| - |1\rangle\langle 1|) + H.c. \quad (2)$$

where δ is the average one-photon detuning, Δ is the two-photon detuning and $H.c.$ is the Hermitian conjugate. We phenomenologically introduce the excited state decay rate Γ with equal rates to $|1\rangle$ and $|2\rangle$, and ground state population difference (coherence) decay rate γ_1 (γ_2). The Rabi frequency is assumed to be much smaller than Γ , so the excited state can be eliminated and the optical coherence adiabatically follows the ground states.

Since for typical diode lasers, the effect of phase noise on atom-light interaction dominates over that of the intrinsic intensity noise, we neglect the latter in our model. The PN converted IN, δI_1 and δI_2 are obtained for cross-correlation calculation: $g^{(2)}(0) = \frac{\langle \delta I_1 \delta I_2 \rangle}{\sqrt{\langle (\delta I_1)^2 \rangle \langle (\delta I_2)^2 \rangle}}$ where $\langle \rangle$ is the time average. In the optically thin regime, intensity fluctuations are proportional to coherence fluctuations of atoms, which gives the zero time lag cross correlation of the INs [9]

$$g^{(2)}(0) = \frac{\langle \text{Im}(\delta \rho_{31}(t)) \text{Im}(\delta \rho_{32}(t)) \rangle}{\sqrt{\langle (\text{Im}(\delta \rho_{31}(t)))^2 \rangle \langle (\text{Im}(\delta \rho_{32}(t)))^2 \rangle}} \quad (3)$$

where ρ_{ij} is the element of the atomic density matrix, and $\langle \rangle$ is the ensemble average.

The average terms in Eq. (3) can be rewritten as $\langle \text{Im}(\delta \rho_{ij}) \text{Im}(\delta \rho_{kl}) \rangle = \frac{1}{4}[\langle \rho_{ij}, \rho_{lk} \rangle + \langle \rho_{ji}, \rho_{kl} \rangle - \langle \rho_{ij}, \rho_{kl} \rangle - \langle \rho_{ji}, \rho_{lk} \rangle]$, where $\langle A, B \rangle = \langle AB \rangle - \langle A \rangle \langle B \rangle$. We then define a vector composed of the independent elements of the density matrix $\mathbf{u} = (\rho_{bb}, \rho_{cc}, \rho_{ab}, \rho_{ba}, \rho_{ac}, \rho_{ca}, \rho_{bc}, \rho_{cb})^T$ where $\rho_{aa} + \rho_{bb} + \rho_{cc} = 1$ was applied. Following the perturbation treatment in [19] using Eq. (1), the averaged master equation can be written as

$$\partial_t \langle \mathbf{u} \rangle = [\mathbf{A}_0 + 2D\mathbf{B}^2] \langle \mathbf{u} \rangle + \mathbf{b} \quad (4)$$




FIG. 2: (Color online) Numerical calculation results. (a-b) $g^{(2)}(0)$ spectra at various Rabi frequencies for laser width $D = 1$ MHz and $D = 81$ MHz. (c) $g^{(2)}(0)$ width vs. laser power for different laser width. (d) Saturated $g^{(2)}(0)$ width vs. laser linewidth. Simulation parameters: ground state coherence decay rate $\gamma_2 = \pi \times 78$ kHz, excited state decay rate $\Gamma = 2\pi \times 500$ MHz.

where $\mathbf{A}_0$ and $\mathbf{b}$ contain Rabi frequency, decay rates and detunings, and $\mathbf{B}$ is a constant diagonal matrix.

To compute the motion of $\langle \text{Im}(\delta \rho_{ij}) \text{Im}(\delta \rho_{kl}) \rangle$, we define a matrix $\mathbf{U} = \mathbf{u}^T \mathbf{u} - \langle \mathbf{u} \rangle^T \langle \mathbf{u} \rangle$, and perform the same stochastic average and projection as for Eq. (4), and obtain

$$\partial_t \langle \mathbf{U} \rangle = [\tilde{\mathbf{A}}_0 + 2D\tilde{\mathbf{B}}_1^2] \langle \mathbf{U} \rangle - 2D(\tilde{\mathbf{B}}_2 - \tilde{\mathbf{B}}_1^2) \langle \tilde{\mathbf{U}} \rangle. \quad (5)$$

where $\tilde{\mathbf{A}}_0$ and $\tilde{\mathbf{B}}_1$ are of similar origins with $\mathbf{A}_0$ and $\mathbf{B}$ in Eq. (4) and $\tilde{\mathbf{B}}_2$ is a constant matrix arising in the projection from $\langle \mathbf{u} \rangle^T \langle \mathbf{u} \rangle$ to $\langle \tilde{\mathbf{U}} \rangle$. Since the slowly varying envelope of $\langle \mathbf{U} \rangle$ is of our interest, the stationary solution of Eq. (5) is taken to compute the cross correlation. For a two level-system, analytical solution for Eq. (5) has been derived, providing a clear picture of PN to IN conversion [20–22]; however, explicit forms of the solution for a three-level system are difficult to obtain due to the much more cumbersome algebra.

We have performed numerical calculation of Eq. (5) and the main results are summarized in Fig. 2. Fig. 2 (a-b) show the lineshape of $g^{(2)}(0)$ spectra at several Rabi frequencies for a small and a large laser linewidth respectively, manifesting as a central peak superimposed on a broader dip-shaped structure resulted from the power broadened wing of $g^{(2)}(0)$ spectrum [8]. At small optical pumping rate, complete anticorrelation cannot be reached because of weak ground state coherence, and the central peak merges with the wing whose linewidth is on the scale of the optical pumping rate. At higher laser

power, the outer wing structure is completely separated from the central narrow peak.

The width of $g^{(2)}(0)$ is defined to be the full width at half of the central peak's amplitude. Fig. 2(c) shows the linewidth vs. Rabi frequency for different laser linewidth. When the laser linewidth is larger, the central peak itself is broader because the laser frequency samples through a wider range, "seeing" more of the parallel slopes; but on the other hand, the optical pumping rate $\Gamma'_p = \frac{\Omega^2}{2(\Gamma+2D)}$ and correspondingly the wing width is smaller. These two effects act together to merge the center and wing structure of the $g^{(2)}(0)$ spectrum, and as a consequence, the $g^{(2)}(0)$ width increases with Rabi frequency since the outer wing's linewidth increases with Rabi frequency. For smaller laser linewidth, the central peak separates better from the outer wing, and so the $g^{(2)}(0)$ width behavior is dominated by the central peak whose width decreases with increasing Rabi because the offset (see Fig. 1) is larger for higher laser power, making the anticorrelation appear sooner. For all laser widths, the $g^{(2)}(0)$ width is saturated at high Rabi frequency because full anticorrelation is reached. The saturated $g^{(2)}(0)$ width increases with laser linewidth but all remain below the lifetime limited width $2\gamma_2$ (Fig. 2 (c)).

To test the above theoretical results, experiments are carried out using a ^{87}Rb enriched vacuum vapor cell under CPT configuration with ground states being Zeeman sublevels of $5^2S_{1/2}$, $F = 2$ and excited states being Zeeman sublevels of $5^2P_{1/2}$, $F' = 1$. The effective ground state coherence decay rate in our system is determined by the transit time of the atoms crossing the laser beam. Within a three-layer magnetic shield, the vapor cell was placed in a solenoid which produces a homogenous magnetic field for two-photon detuning adjustment. The laser beam was linearly polarized before entering the cell, and its right and left circular polarized components acted as E_1 and E_2 in Fig. 1 and are separately detected at the cell output by amplified photo-detectors. ac signals were acquired at a local computer from an oscilloscope and the $g^{(2)}(0)$ value was computed offline.

We first used an external cavity diode laser with linewidth < 1 MHz. The cell temperature was 52°C . EIT FWHM vs. laser power was measured as shown in Fig. 3(b) (dark dots). We extrapolated the transit time limited width by fitting the data with polynomials and extending the fitting result to zero power. The obtained full width is 75 kHz, close to the transit time limited width estimated from the measured laser beam diameter of 2.8 mm. The $g^{(2)}(0)$ vs. Δ spectra at different laser power is shown in Fig. 3(a). The FWHM of the central peaks of $g^{(2)}(0)$ vs. laser power are plotted along with EIT linewidth in Fig. 3(b). The narrowest linewidth of $g^{(2)}(0)$ observed is about 1/3 of the transit linewidth of EIT. At small laser power, the $g^{(2)}(0)$ spectrum can not achieve perfect anticorrelation because of the weak ground state coherence. Besides, electronics noise from the photo-detectors makes INs less correlated at $\Delta = 0$. The trend of the $g^{(2)}(0)$ width vs. laser power is consis-




FIG. 3: (Color online) Experimental results using a laser with linewidth about 1 MHz. (a) $g^{(2)}(0)$ spectra at various laser powers. (b) EIT and $g^{(2)}(0)$ linewidth vs. laser power. In the $g^{(2)}(0)$ width curve, the line is to guide the eye. The dashed line in (b) is the fitted zero power EIT width with polynomial.




FIG. 4: (Color online) Experimental results using a laser with linewidth about 80 MHz. (a) $g^{(2)}(0)$ spectra at various laser powers. (b) EIT and $g^{(2)}(0)$ linewidth vs. laser power. In the $g^{(2)}(0)$ width curve, the line is to guide the eye. The dashed line in (b) is the fitted zero power EIT width with polynomial.

tent with our theoretical prediction for small laser width, except for slight broadening at large laser power shown in the inset in Fig. 3(b). We attribute this broadening to increased influence of laser intensity noise on $g^{(2)}(0)$ at higher laser power.

In a second experiment, we used a vertical cavity surface emitting laser (VCSEL) with a linewidth of ~ 80 MHz. The vapor cell temperature was at 40°C to avoid excess absorption of the relatively weak laser beam (maximal power was $250 \mu\text{W}$). The $g^{(2)}(0)$ spectra and the width along with EIT width are obtained at different laser power shown in Fig. 4. The extrapolated transit width of EIT is 61 kHz, close to the transit width estimated from the laser diameter of 2.5 mm. $g^{(2)}(0)$ linewidth is also below the transit width. The linewidth increase with laser power is consistent with theory prediction for larger laser linewidth.

In both experiments, measured $g^{(2)}(0)$ width is larger than predicted. Several factors neglected in the model contribute to the discrepancy: (a) intrinsic intensity noise

from the laser in E_1 and E_2 are correlated and hence reduces anticorrelation; (b) electronics noise in the photo detector and data acquisition system, and eventually laser shot noise (negligible compared to other noise in our system) are not correlated and bring the $g^{(2)}(0)$ towards zero; (c) Our system is not strictly in the optically thin regime, and studies show that $g^{(2)}(0)$ linewidth increases with optical depth [23]. These noise sources and complications prevent the linewidth from approaching zero as predicted in our simple model.

We note that the sharp transition from correlation to anticorrelation here has an inherent connection with the predicted sudden appearance of entanglement between collective optical modes when Raman detuning deviates from zero [24]. In ref [24], a phase sensitive EIT based on a double Λ system is considered, where each Λ couples two optical fields, and the two fields coupling to the same ground state form a quantum beat and hence can be viewed as a collective field. When the Raman detuning is zero, atoms are transparent to light and cannot induce quantum correlations to the lasers; once the Raman detuning deviates from zero, the sharp dispersion of the ground state coherence drastically modifies the statistics of the fields, and entanglement occurs between

any two fields from different collective modes. In brief, the connection lies in that, in our case, atom-light interaction induces anticorrelation when exact EIT resonance condition is broken, and in [24], it introduces quantum correlation between optical fields.

In conclusion, we theoretically and experimentally studied the laser intensity noise correlations generated in atom-light interactions with a phase diffused laser in a CPT system, and found that the transition from correlation to anticorrelation can be very sharp due to the ground state coherence, with a linewidth below the lifetime-limited width and theoretically approaching zero at small laser width. This result is relevant to the realization of recent proposals of creating quantum correlations between two bright optical fields and also among atomic spins using CPT, and may be also used for spectroscopy.

We thank Michael Crescimanno, Xiangming Hu, Liang Jiang, Jianming Wen, Saijun Wu and Michael Hohensee for stimulating discussions. This work is funded by the NBRPC (973 Program Grants No. 2012CB921604 and No. 2011CB921604), NNSFC (Grant No. 10904018) and Fudan University. Correspondence should be sent to yxiao@fudan.edu.cn.

-
- [1] M. Fleischhauer, A. Imamoglu, J. P. Marangos, *Rev. Mod. Phys.* **77**, 633 (2005).
 - [2] J. Vanier, *Appl. Phys. B* **81**, 421 (2007).
 - [3] A. Dantan, J. Cviklinski, E. Giacobino, M. Pinard, *Phys. Rev. Lett.* **97**, 023605 (2006).
 - [4] A. Sinatra, *Phys. Rev. Lett.* **97**, 253601 (2006).
 - [5] P. Barberis-Blostein and M. Bienert, *Phys. Rev. Lett.* **98**, 033602 (2007).
 - [6] A. Godone, S. Micalizio, and F. Levi, *Phys. Rev. A* **66**, 063807 (2002).
 - [7] M. Martinelli, P. Valente, H. Failache, D. Felinto, L. S. Cruz, P. Nussenzeig and A. Lezama, *Phys. Rev. A* **69**, 043809 (2004).
 - [8] Y. Xiao, T. Wang, M. Baryakhtar, M. V. Camp, M. Crescimanno, M. Hohensee, L. Jiang, D. F. Phillips, M. D. Lukin, S. F. Yelin, and R. L. Walsworth, *Phys. Rev. A* **80**, 041805(R) (2009).
 - [9] V. A. Sautenkov, Y. V. Rostovtsev, and M. O. Scully, *Phys. Rev. A* **72**, 065801 (2005).
 - [10] L. Feng, P. Li, L. Jiang, J. Wen, and Y. Xiao, *Phys. Rev. Lett.* **109**, 233006 (2012).
 - [11] J. C. Camparo, J. G. Coffer, *Phys. Rev. A* **59**, 728(1999).
 - [12] S. A. Crooker, D. G. Rickel, A. V. Balatsky, and D. L. Smith, *Nature (London)* **431**, 49 (2004).
 - [13] N. Muller and A. Jerschow, *Proc. Natl. Acad. Sci. U.S.A.* **103**, 6790 (2006).
 - [14] M. Bahoura, A. Clairon, *Opt. Lett.*, **26**, 926 (2001).
 - [15] The offset only exists when the ground state population difference decay rate γ_1 is nonzero. However, the condition of anticorrelations is not the offset, but a nonzero imaginary part of ρ_{12} .
 - [16] C. W. Gardiner, *Quantum Noise* (Springer-Verlag, Berlin, 2000).
 - [17] M. O. Scully, and M. S. Zubairy, *Quantum Optics* (Cambridge University Press, Cambridge, 1997).
 - [18] A. Godone, F. Levi, and S. Micalizio, *Coherent Population Trapping Maser* (C. L. U. T, Torio, 2002).
 - [19] N. G. Van Kampen, *Stochastic Processes in Physics and Chemistry* (Elsevier, Singapore, 1997).
 - [20] P. Zoller, *Phys. Rev. A* **19**, 1151 (1979).
 - [21] T. Haslwanter, H. Ritsch, J. Cooper, and P. Zoller, *Phys. Rev. A* **38**, 5652 (1988).
 - [22] H. Ritsch and P. Zoller, *Phys. Rev. A* **41**, 2654 (1990).
 - [23] P. Li, L. Feng *et.al.*, to be published.
 - [24] F. Wang, X. Hu, W. Shi, Y. Zhu, *Phys. Rev. A* **81**, 033836 (2010).